


IMPORTANT SAFETY INFORMATION

Amgen Inc.
One Amgen Center Drive
Thousand Oaks, CA 91320-1779
www.amgen.com
© 2020 Amgen Inc. All rights reserved.
USA-162-82567 11/20

[Privacy Statement](#) | [Terms of Use](#) | [Contact Us](#)

Attn: Subscriptions
2 Clarke Drive, Ste 100
Cranbury, NJ 08512
USA
[Privacy Policy](#)



For patients requiring invasive dental procedures, clinical judgment should guide the management plan of each patient. Patients who are suspected of having or who develop ONJ should receive care by a dentist or an oral surgeon. Extensive dental surgery to treat ONJ may exacerbate the condition. Discontinuation of Prolia® should be considered based on individual benefit-risk assessment.

Atypical Femoral Fractures: Atypical low-energy, or low trauma fractures of the shaft have been reported in patients receiving Prolia®. Causality has not been established as these fractures also occur in osteoporotic patients who have not been treated with antiresorptive agents.

During Prolia® treatment, patients should be advised to report new or unusual thigh, hip, or groin pain. Any patient who presents with thigh or groin pain should be evaluated to rule out an incomplete femur fracture. Interruption of Prolia® therapy should be considered, pending a risk/benefit assessment, on an individual basis.

Multiple Vertebral Fractures (MVF) Following Discontinuation of Prolia® Treatment: Following discontinuation of Prolia® treatment, fracture risk increases, including the risk of multiple vertebral fractures. New vertebral fractures occurred as early as 7 months (on average 19 months) after the last dose of Prolia®. Prior vertebral fracture was a predictor of multiple vertebral fractures after Prolia® discontinuation. Evaluate an individual's benefit/risk before initiating treatment with Prolia®. If Prolia® treatment is discontinued, patients should be transitioned to an alternative antiresorptive therapy.

Serious Infections: In a clinical trial (N = 7808), serious infections leading to hospitalization were reported more frequently in the Prolia® group than in the placebo group. Serious skin infections, as well as infections of the abdomen, urinary tract and ear, were more frequent in patients treated with Prolia®.

Endocarditis was also reported more frequently in Prolia®-treated patients. The incidence of opportunistic infections and the overall incidence of infections were similar between the treatment groups. Advise patients to seek prompt medical attention if they develop signs or symptoms of severe infection, including cellulitis.

Patients on concomitant immunosuppressant agents and with impaired immune systems may be at increased risk for serious infections. In patients who develop serious infections while on Prolia®, prescribers should assess the need for continued Prolia® therapy.

Dermatologic Adverse Reactions: Epidermal and dermal adverse events such as dermatitis, eczema and rashes occurred at a significantly higher rate with Prolia® compared to placebo. Most of these events were not specific to the injection site. Consider discontinuing Prolia® if severe symptoms develop.

Musculoskeletal Pain: Severe and occasionally incapacitating bone, joint, and/or muscle pain has been reported in patients taking Prolia®. Consider discontinuing use if severe symptoms develop.

Suppression of Bone Turnover: Prolia® resulted in significant suppression of bone remodeling as evidenced by markers of bone turnover and bone histomorphometry. The significance of these findings and the effect of long-term treatment are unknown. Monitor patients for consequences, including ONJ, atypical fractures, and delayed fracture healing.

Adverse Reactions: The most common adverse reactions (>5% and more common than placebo) are back pain, pain in extremity, musculoskeletal pain, hypercholesterolemia, and cystitis. Pancreatitis has been reported with Prolia®.

The overall incidence of new malignancies was 4.3% in the placebo group and 4.8% in the Prolia® group. A causal relationship to drug exposure has not been established. Denosumab is a human monoclonal antibody. As with all therapeutic proteins, there is potential for immunogenicity.

Please see Prolia® full Prescribing Information, including Medication Guide.

Reference: 1. Data on file, Amgen; 2020.

Amgen Inc.
One Amgen Center Drive
Thousand Oaks, CA 91320-1799
www.amgen.com
© 2020 Amgen Inc. All rights reserved.
USA-162-82567 11/20

AMGEN

Privacy Statement | Terms of Use | Contact Us

You received this e-mail because you are a customer of Managed Healthcare Executive, a MJH Life Sciences publication. You are on the mailing list as SUSERS@DOMAIN

To ensure delivery to your Inbox, please add us to your address book.

From time to time, we will inform customers of latest offers and new products from relevant business partners. If you do not wish to receive this type of information, [click here](#).

Contact us via email at rmhinfo@rmhgroup.com or contact us by mail at

MJH Life Sciences
Attn: Subscriptions
2 Clarke Drive, Ste 100
Cranbury, NJ 08512
USA
[Privacy Policy](#)